

University of Groningen

Novel insights into FXR actions in liver and intestine

Boesjes, Marije

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2015

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Boesjes, M. (2015). *Novel insights into FXR actions in liver and intestine*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

General introduction, scope and outline



GENERAL INTRODUCTION

Bile salts

Bile formation and bile salt synthesis

The biliary pathway is one of the major routes through which endogenous waste products (*e.g.* bilirubin), drugs and other toxic compounds are eliminated from the body and also represents a main route for removal of excess cholesterol. Hence, formation of bile is an important physiological function of the liver. Primary bile is formed by hepatocytes and active secretion of bile salts (BS) into the bile canaliculi provides the main driving force for bile formation. Upon release in the small intestine, BS promote the absorption of dietary fats and fat-soluble vitamins. BS are synthesized from cholesterol in the hepatocytes by well-characterized biosynthetic pathways. The classical (neutral) pathway starts with 7 α -hydroxylation of cholesterol by cholesterol 7 α -hydroxylase (CYP7A1) while the alternative (acidic) pathway is initiated by sterol 27-hydroxylase (CYP27A1)¹. Newly synthesized, so-called primary BS, are cholate (CA) and chenodeoxycholate (CDCA) in humans. In rodents, the relatively hydrophobic CDCA molecules are rapidly hydroxylated to form the more hydrophilic alpha- and beta-muricholates (α -MCA and β -MCA) (Fig. 1). Sterol 12 α -hydroxylase (CYP8B1) activity is required for the biosynthesis of CA and determines the ratio CA to CDCA and, therefore, the hydrophobicity of the BS




Figure 1. Bile salt structure

Structure of major naturally occurring bile salts. The primary bile salts in humans are CA; cholate and CDCA; chenodeoxycholate and the secondary bile salts in humans are DCA; deoxycholate, LCA; lithocholate and UDCA; ursodeoxycholate. The primary bile salts in rodents are CA, α MCA; alpha-muricholate and β MCA; beta-muricholate and the secondary bile salts in rodents are HCA; hyocholate, MDCA; murideoxycholate, ω MCA; omega-muricholate and HDCA; hyodeoxycholate.

pool ². The diversity of BS species in the BS pool is expanded by conjugation with amino acids to either taurine in rodents or glycine in humans in the liver and by the actions of intestinal bacteria that, *via* deconjugation, oxidation of hydroxyl groups and dehydroxylation, form the secondary BS deoxycholate (DCA) and lithocholate (LCA) in humans, and hyocholate (HCA), hyodeoxycholate (HDCA), murideoxycholate (MDCA), omega-muricholate (ω -MCA) and ursodeoxycholate (UDCA) in rodents (Fig. 1) ³. These processes collectively creates a BS pool with specific physiochemical properties.

Enterohepatic circulation of bile salts

BS are maintained efficiently in the enterohepatic circulation (Fig. 2); 95% of the BS secreted by the liver is reabsorbed in the small intestine and only 5% is lost in the feces. The liver compensates for this loss by *de novo* BS synthesis. In the intestine, the reabsorption of bile salts is regulated through the expression of four important proteins, *i.e.*, apical sodium-dependent bile acid transporter (SLC10A2/IBAT/ASBT) ⁴, intestinal bile acid-binding protein (IBABP/FABP6) ⁵ and the heterodimer organic solute transporters alpha (SLC51A /OSTA) and beta (SLC51B /OSTB) ⁶, that are collectively responsible for the transport of BS from the intestinal lumen to the portal system. Na⁺-taurocholate cotransporting polypeptide (SLC10A1/NTCP) ⁷ and organic anion-transporting peptides (OATPs/SLCO superfamily) ⁸ are the major BS transporters at the basolateral membranes of hepatocytes for the uptake of BS and organic solutes from the portal system to the liver. Reabsorbed BS will be conjugated again and subsequently transported into bile by the bile salt export pump (BSEP/ABCB11) ⁹.

The Farnesoid X receptor

Nuclear receptors

Nuclear receptors are a class of proteins that are able to sense the cellular presence of nutrients, steroid hormones and certain other molecules like vitamins and BS. Upon activation, these receptors interact with other proteins to regulate the expression of target genes, thereby controlling a vast variety of cellular processes. Examples of nuclear receptors, their endogenous ligands and their main functions are the peroxisome proliferator-activated receptors (NR1C1/PPAR α ; NR1C2/PPAR β / δ ;




Figure 2. The enterohepatic circulation of bile salts

Bile salts (BS) are synthesized from cholesterol in hepatocytes and are actively secreted into the bile canaliculi that drain into the common bile duct. BS are stored in the gallbladder and upon ingestion of a meal the content is released into the intestinal lumen. In the ileum, BS are taken up via ASBT into the ileocytes and BS activate FXR which, together with RXR, induce the expression of Fgf15. FGF15 is released into the circulation and suppresses the hepatic BS synthesis by inhibiting Cyp7a1 expression. BS returning to the liver after reabsorption in the intestine also contribute to the negative feedback loop by suppression of Cyp7a1 via hepatic FXR activation. FXR; Farnesoid X Receptor, RXR; Retinoid X Receptor; CYP7A1; cholesterol 7 α -hydroxylase, FGFR4; fibroblast growth factor receptor 4, FGF15; fibroblast growth factor 15, ASBT; apical sodium-dependent bile acid transporter, IBABP; intestinal bile acid-binding protein, OST α/β ; organic solute transporters alpha/beta, NTCP; Na⁺-taurocholate cotransporting polypeptide, OATPs; organic anion-transporting peptides.

NR1C3/PPAR γ) that upon activation by free fatty acids function in control of cellular differentiation and lipid metabolism ¹⁰, Liver X Receptors (NR1H3/LXR α ; NR1H2/LXR β) that upon activation by oxysterols modulate cholesterol metabolism ¹¹ and Farnesoid X Receptor (NR1H4/FXR α , from now on referred to as FXR ¹²⁻¹⁴ that upon activation by BS regulates steps in BS, lipid and glucose metabolism. The classical view of structural organization divided a nuclear receptor protein into 5 domains; the N-terminal transactivation independent regulatory domain (AF1), the DNA-binding domain (DBD), a flexible hinge region (H), a ligand-binding domain (LBD) and a C-terminal transactivation dependent domain (AF2) ¹⁵. Nuclear receptors can bind DNA as homodimers, heterodimers or monomers ¹⁶. Nuclear receptors have to ability to directly bind to DNA to regulate transcription or bind DNA as heterodimers with the common partner Retinoid X Receptor (NR2B1/RXR α ; NR2B2/RXR β) ¹⁷. Upon binding to specific DNA sequences in the cell nucleus, nuclear receptors recruit transcriptional co-regulators, including transcription factors, co-activators, co-repressors and the RNA machinery itself, which all effect transcription ¹⁸.

Bile salt-activated receptor FXR


BS regulate their own synthesis in feed-back mechanisms involving FXR in the intestine and liver, the two organs in which FXR is highly expressed. FXR is the major BS responsive ligand-activated transcription factor and is involved in control of BS homeostasis. Intestinal BS-activated FXR induces the expression of fibroblast growth factor 19 (FGF19, and its mouse homolog Fgf15), which is secreted into portal blood to down-regulate hepatic BS synthesis *via* fibroblast growth factor receptor 4 (FGFR4) and B-Klotho signaling in the liver ^{19,20}. Activation of the FGFR4/B-Klotho (KLB) complex stimulates the c-Jun N-terminal kinase (JNK) pathway, suppressing hepatic *Cyp7a1* expression that catalyzes the rate-controlling step in the BS synthesis (Fig. 2). Hepatic BS levels are maintained by the control of uptake, metabolism, synthesis and excretion of BS. Activated hepatic FXR reduces *Cyp7a1* expression, in a short heterodimer partner (NR0B2/SHP)- and liver receptor homolog-1 (NR5A2/LRH1)-dependent manner ^{21,22}. Also *Cyp8b1*, determining the CA to CDCA ratio, and *Cyp27a1*, regulating the alternative BS biosynthetic pathway, are regulated by BS-activated FXR, in a SHP and hepatocyte nuclear factor-4 alpha (NR2A1/HNF4A)-dependent manner ²³⁻²⁶. FXR can be activated by its endogenous ligands, *i.e.*, BS, with species-specific potencies. The relatively hydrophobic BS appear to be the most potent ones; CDCA>DCA>LCA>CA ¹³. However, also several

synthetic compounds have been developed the past years, such as the BS derivative 6-ethyl-chenodeoxycholic acid (INT747) ²⁷, but also non-steroid compounds such as GW4064 ²⁸, WAY-362450 (XL335) ²⁹, PX20350 and PX20606 ³⁰.


FXR isoforms

FXR has four different splice variants in both humans and rodents^{31,32}. These isoforms arise through alternative splicing of exon 5 and the use of distinct promoters that initiate transcription from either exon 1 or exon 3 (Fig. 3A and B). The different promoters of the *FXR* gene regulate the expression of FXRa1,2 or FXRa3,4 transcripts. FXRa1 and FXRa3 transcripts contain four amino acids (MYTG) in the hinge domain,

A *Fxr/Nr1h4* gene structure



B FXR isoforms



C Amino acid sequence of N-terminal regulatory domains

FXRa1/2 - - MNLIGHS- - - HLQATDEFSLS - - - - - - - - ESFLG
FXRα3/4 MVMOFOGLNPIQISLHSHRLSGFVPEGMSVKPAKG

Figure 3. The Farnesoid X Receptor

Schematic representation of the murine *Fxr/Nr1h4* gene (A) and FXR isoforms (B) with two distinct N-terminal regulatory domains (C). Two alternative promoters give rise to FXR isoforms FXRa1/2 and FXRa3/4 and a 12 base pair alternative splice donor site, which determines the difference between FXRa1/3 and FXRa2/4. AF1; transactivation independent domain, AF2; transactivation dependent domain, DBD; DNA binding domain; LBD; ligand binding domain, H; hinge region, MYTG; 4-amino acids insertion. transcription start site, * translation start site.

adjacent to the DNA-binding domain (DBD), due to alternative splicing. The FXR isoforms show differences in spatial and temporal expression, as well as transcriptional activity^{33,34}. In mice all four isoforms are abundantly expressed in liver, while FXR α 3 and α 4 are abundantly expressed in ileum, moderately in kidney and at low levels in stomach, duodenum, and jejunum. FXR α 1 and α 2 are moderately expressed in ileum and adrenal gland³². The difference between FXR α 2 and FXR α 4 lies within the N-terminal part of the protein, *i.e.*, resulting in a 37 amino acid extension of FXR α 4 (Fig. 3C). This difference can cause conformational changes in the FXR protein itself and, thereby, influence the transcriptional activity and/or transactivation/repression capacity on target genes. A dozen transcriptional cofactors have been described to influence FXR transactivation³⁵, but actions specific for the FXR isoforms have not been described. The intriguing question remains why FXR isoforms exist and what their specific physiological roles in different tissues actually are.

The expression of FXR itself is tightly regulated, there are several cofactors that play a role in FXR-mediated transcription. PGC-1 α , a well-characterized coactivator for PPAR α and the master integrator of cellular energy metabolism, coactivates FXR-target gene promoter activity *in vitro*³⁶⁻³⁸. Coactivators such as SMARCA4, SIRT1, EP300, CARM1, PRMT1 and ASCOM are recruited to FXR target genes, which results in histone modification or chromatin remodeling and increased gene transcription³⁹. Interestingly, some of these cofactors, including EP300 and SIRT1, acetylate and deacetylate not only histones at FXR target genes but also FXR itself³⁹. Molecular regulation of FXR activity also involves post-transcriptional phosphorylation, sumoylation and ubiquitination³⁹. In addition, FXR expression levels vary with the nutritional status and appear to be tightly regulated by the metabolic status, indirectly supporting a role for this nuclear receptor in metabolic diseases^{37,40-42}.

The relative abundance of hepatic FXR α 3/4 was increased in children suffering from progressive familial cholestasis type I⁴³. Also in cholangiocarcinomas⁴⁴ and colorectal adenocarcinoma⁴⁵ the relative abundance of hepatic and intestinal FXR α 3/4 was shown to be increased. Apparently, disease states can be associated with a switch in FXR isoform expression in the liver and intestine which may influence outcome of the disease state progress. Recently, Vaquero *et al.*⁴⁶ reported that activation of human FXR depends on the pattern of FXR isoform expression and the BS composition. They showed that cell-specific pattern of FXR isoforms determine the overall tissue sensitivity to FXR agonists and may be involved in the differential response of FXR target genes to FXR activation.

Bile salt-activated FXR as metabolic regulator

BS-activated FXR turned out to be involved in control of lipoprotein, glucose and energy metabolism ⁴⁷. A number of these findings became evident from studies in FXR-deficient mice. FXR-deficiency was associated with reduced expression of the gluconeogenic enzyme PECK ^{48,49}, impaired glucose tolerance, insulin resistance ^{50,51}, hypertriglyceridemia and increased VLDL production ^{52,53}. Furthermore, administration of the natural FXR ligands, BS, resulted in increased energy expenditure in brown adipose tissue of mice fed a high fat diet ⁵⁴. Clinical observations showed that the treatment of patients with BS sequestrants, *i.e.* removal of natural ligands, also results in increased plasma triglyceride (TG) levels ^{55,56}, while CDCA-treatment had the opposite effect ^{55,57,58}. The generation of synthetic FXR compounds provided new insights into the metabolic actions of FXR. *In vivo* treatment with the synthetic FXR agonist GW4064 significantly improved hypercholesterolemia, hypertriglyceridemia and hyperglycemia in mouse models of insulin resistance ^{48,50,51}. Furthermore, the synthetic FXR agonist PX20350/20606 induced HDL-mediated transhepatic cholesterol efflux in mice and monkeys and prevented atherosclerosis in a susceptible mouse model ^{30,59}. Note that BS are also able to activate the transmembrane receptor TGR5, also known as G-protein coupled BS receptor 1 (GPBAR1) and membrane-type BS receptor ⁶⁰. BS-activated TGR5 is involved in glucose metabolism, energy expenditure, immune response and liver function ⁶¹⁻⁶³.

SCOPE AND OUTLINE

FXR is considered a promising target for novel therapies aimed at treatment of metabolic derangements associated with obesity, *i.e.* type II diabetes, non-alcoholic fatty liver disease and atherosclerosis. Activated FXR plays key roles in regulating metabolism of BS, lipids and glucose. The fact that four different FXR isoforms exist, and that these are conserved in rodent and human genomes argues for a functional role of these isoforms. The FXR isoform specific function regarding metabolic regulation is currently unknown and requires clarification.

Altogether, insight into tissue-specific actions of the FXR isoforms as well as in the consequences of alterations in BS pool size and composition are required to define optimal strategies to treat and prevent components of the metabolic syndrome. Using innovative mouse models and by modulating the activity of FXR,

we have evaluated their role in control of BS and cholesterol metabolism. **Chapter 2** reviews the metabolic effects of BS in the gut in health and disease. In **chapter 3**, we investigated the tissue-specific roles of FXR α 2 and FXR α 4, the transcriptionally most active splice variants of FXR, on BS biosynthesis and lipoprotein metabolism in the liver. We show distinct roles for FXR α 2 and FXR α 4 in the different branches of the BS biosynthetic pathway. Furthermore, mice with hepatic FXR α 2 expression showed a relatively hydrophilic BS pool and increased their cholesterol turnover compared to FXR α 4 mice. **Chapter 4** focusses on the pharmacological activation of both LXR and FXR, to overcome the hepatic triglyceride (TG) accumulation caused by pharmacological activation of LXR alone. We show that concomitant activation of both nuclear receptors ameliorated hepatic steatosis, possibly *via* increased TG-hydrolysis induced by pharmacological FXR activation. We also assessed the contribution of intestinal FXR on BS and cholesterol homeostasis, which is described in **chapter 5**. Our data revealed that intestinal FXR is capable of mediating removal of large quantities of cholesterol from the body. Stimulation of the TICE pathway was shown to contribute most to the increased cholesterol disposal in response to FXR activation. Furthermore, our data pinpoint intestinal FXR as a transcription factor impacting the hydrophobicity of the BS pool thereby governing the rate of transintestinal cholesterol excretion.

REFERENCES

1. Russell DW. The enzymes, regulation, and genetics of bile acid synthesis. *Annu Rev Biochem* 2003;72:137-174.
2. Vlahcevic ZR, Eggertsen G, Bjorkhem I, Hylemon PB, Redford K, Pandak WM. Regulation of sterol 12 α -hydroxylase and cholic acid biosynthesis in the rat. *Gastroenterology* 2000;118:599-607.
3. Kuipers F, Bloks VW, Groen AK. Beyond intestinal soap-bile acids in metabolic control. *Nat Rev Endocrinol* 2014;10:488-498.
4. Shneider BL. Intestinal bile acid transport: biology, physiology, and pathophysiology. *J Pediatr Gastroenterol Nutr* 2001;32:407-417.
5. Coppola CP, Gosche JR, Arrese M, et al. Molecular analysis of the adaptive response of intestinal bile acid transport after ileal resection in the rat. *Gastroenterology* 1998;115:1172-1178.
6. Dawson PA, Hubbert M, Haywood J, et al. The heteromeric organic solute transporter alpha-beta, Ostalpha-Ostbeta, is an ileal basolateral bile acid transporter. *J Biol Chem* 2005;280:6960-6968.
7. Stieger B, Hagenbuch B, Landmann L, Hochli M, Schroeder A, Meier PJ. In situ localization of the hepatocytic Na⁺/Taurocholate cotransporting polypeptide in rat liver. *Gastroenterology* 1994;107:1781-1787.
8. Hagenbuch B, Meier PJ. Organic anion transporting polypeptides of the OATP/ SLC21 family: phylogenetic classification as OATP/ SLCO superfamily, new nomenclature and molecular/ functional properties. *Pflugers Arch* 2004;447:653-665.
9. Gerloff T, Stieger B, Hagenbuch B, et al. The sister of P-glycoprotein represents the canalicular bile salt export pump of mammalian liver. *J Biol Chem* 1998;273:10046-10050.
10. Monsalve FA, Pyarasani RD, Delgado-Lopez F, Moore-Carrasco R. Peroxisome proliferator-activated receptor targets for the treatment of metabolic diseases. *Mediators Inflamm* 2013;2013:549627.
11. Oosterveer MH, Grefhorst A, Groen AK, Kuipers F. The liver X receptor: Control of cellular lipid homeostasis and beyond Implications for drug design. *Prog Lipid Res* 2010;.
12. Makishima M, Okamoto AY, Repa JJ, et al. Identification of a nuclear receptor for bile acids. *Science* 1999;284:1362-1365.
13. Parks DJ, Blanchard SG, Bledsoe RK, et al. Bile acids: natural ligands for an orphan nuclear receptor. *Science* 1999;284:1365-1368.
14. Wang H, Chen J, Hollister K, Sowers LC, Forman BM. Endogenous bile acids are ligands for the nuclear receptor FXR/BAR. *Mol Cell* 1999;3:543-553.
15. Mangelsdorf DJ, Thummel C, Beato M, et al. The nuclear receptor superfamily: the second decade. *Cell* 1995;83:835-839.
16. Glass CK. Differential recognition of target genes by nuclear receptor monomers, dimers, and heterodimers. *Endocr Rev* 1994;15:391-407.
17. Mangelsdorf DJ, Evans RM. The RXR heterodimers and orphan receptors. *Cell* 1995;83:841-850.
18. Kinyamu HK, Archer TK. Modifying chromatin to permit steroid hormone receptor-dependent transcription. *Biochim Biophys Acta* 2004;1677:30-45.
19. Holt JA, Luo G, Billin AN, et al. Definition of a novel growth factor-dependent signal cascade for the suppression of bile acid biosynthesis. *Genes Dev* 2003;17:1581-1591.
20. Inagaki T, Choi M, Moschetta A, et al. Fibroblast growth factor 15 functions as an enterohepatic signal to regulate bile acid homeostasis. *Cell Metab* 2005;2:217-225.
21. Goodwin B, Jones SA, Price RR, et al. A regulatory cascade of the nuclear receptors FXR, SHP-1, and LXR-1 represses bile acid biosynthesis. *Mol Cell* 2000;6:517-526.
22. Lu TT, Makishima M, Repa JJ, et al. Molecular basis for feedback regulation of bile acid synthesis by nuclear receptors. *Mol Cell* 2000;6:507-515.
23. Zhang M, Chiang JY. Transcriptional regulation of the human sterol 12 α -hydroxylase gene (CYP8B1): roles of hepatocyte nuclear factor 4 α in mediating bile acid repression. *J Biol Chem* 2001;276:41690-41699.
24. Chen W, Chiang JY. Regulation of human sterol 27-hydroxylase gene (CYP27A1) by bile acids and hepatocyte nuclear factor 4 α (HNF4 α). *Gene* 2003;313:71-82.

25. Garuti R, Croce MA, Piccinini L, Tiozzo R, Bertolini S, Calandra S. Functional analysis of the promoter of human sterol 27-hydroxylase gene in HepG2 cells. *Gene* 2002;283:133-143.
26. Yang Y, Zhang M, Eggertsen G, Chiang JY. On the mechanism of bile acid inhibition of rat sterol 12alpha-hydroxylase gene (CYP8B1) transcription: roles of alpha-fetoprotein transcription factor and hepatocyte nuclear factor 4alpha. *Biochim Biophys Acta* 2002;1583:63-73.
27. Pellicciari R, Fiorucci S, Camaioni E, et al. 6alpha-ethyl-chenodeoxycholic acid (6-ECDCA), a potent and selective FXR agonist endowed with anticholestatic activity. *J Med Chem* 2002;45:3569-3572.
28. Maloney PR, Parks DJ, Haffner CD, et al. Identification of a chemical tool for the orphan nuclear receptor FXR. *J Med Chem* 2000;43:2971-2974.
29. Flatt B, Martin R, Wang TL, et al. Discovery of XL335 (WAY-362450), a highly potent, selective, and orally active agonist of the farnesoid X receptor (FXR). *J Med Chem* 2009;52:904-907.
30. Abel U, Schluter T, Schulz A, et al. Synthesis and pharmacological validation of a novel series of non-steroidal FXR agonists. *Bioorg Med Chem Lett* 2010;20:4911-4917.
31. Huber RM, Murphy K, Miao B, et al. Generation of multiple farnesoid-X-receptor isoforms through the use of alternative promoters. *Gene* 2002;290:35-43.
32. Zhang Y, Kast-Woelbern HR, Edwards PA. Natural structural variants of the nuclear receptor farnesoid X receptor affect transcriptional activation. *J Biol Chem* 2003;278:104-110.
33. Lee FY, Kast-Woelbern HR, Chang J, et al. Alpha-crystallin is a target gene of the farnesoid X-activated receptor in human livers. *J Biol Chem* 2005;280:31792-31800.
34. Lee H, Zhang Y, Lee FY, Nelson SF, Gonzalez FJ, Edwards PA. FXR regulates organic solute transporters alpha and beta in the adrenal gland, kidney, and intestine. *J Lipid Res* 2006;47:201-214.
35. Kemper JK. Regulation of FXR transcriptional activity in health and disease: Emerging roles of FXR cofactors and post-translational modifications. *Biochim Biophys Acta* 2010;.
36. Kanaya E, Shiraki T, Jingami H. The nuclear bile acid receptor FXR is activated by PGC-1alpha in a ligand-dependent manner. *Biochem J* 2004;382:913-921.
37. Zhang Y, Castellani LW, Sinal CJ, Gonzalez FJ, Edwards PA. Peroxisome proliferator-activated receptor-gamma coactivator 1alpha (PGC-1alpha) regulates triglyceride metabolism by activation of the nuclear receptor FXR. *Genes Dev* 2004;18:157-169.
38. Savkur RS, Thomas JS, Bramlett KS, Gao Y, Michael LF, Burris TP. Ligand-dependent coactivation of the human bile acid receptor FXR by the peroxisome proliferator-activated receptor gamma coactivator-1alpha. *J Pharmacol Exp Ther* 2005;312:170-178.
39. Kemper JK. Regulation of FXR transcriptional activity in health and disease: Emerging roles of FXR cofactors and post-translational modifications. *Biochim Biophys Acta* 2011;1812:842-850.
40. Duran-Sandoval D, Mautino G, Martin G, et al. Glucose regulates the expression of the farnesoid X receptor in liver. *Diabetes* 2004;53:890-898.
41. Duran-Sandoval D, Cariou B, Percevault F, et al. The farnesoid X receptor modulates hepatic carbohydrate metabolism during the fasting-refeeding transition. *J Biol Chem* 2005;280:29971-29979.
42. Lefebvre P, Cariou B, Lien F, Kuipers F, Staels B. Role of bile acids and bile acid receptors in metabolic regulation. *Physiol Rev* 2009;89:147-191.
43. Alvarez L, Jara P, Sanchez-Sabate E, et al. Reduced hepatic expression of farnesoid X receptor in hereditary cholestasis associated to mutation in ATP8B1. *Hum Mol Genet* 2004;13:2451-2460.
44. Martinez-Becerra P, Vaquero J, Romero MR, et al. No correlation between the expression of FXR and genes involved in multidrug resistance phenotype of primary liver tumors. *Mol Pharm* 2012;9:1693-1704.
45. Martinez-Becerra P, Monte I, Romero MR, et al. Up-regulation of FXR isoforms is not required for stimulation of the expression of genes involved in the lack of response of colon cancer to chemotherapy. *Pharmacol Res* 2012;66:419-427.
46. Vaquero J, Monte MJ, Dominguez M, Muntane J, Marin JJ. Differential activation of the human farnesoid X receptor depends on the pattern of expressed isoforms and the bile acid pool composition. *Biochem Pharmacol* 2013;86:926-939.
47. Boesjes M, Brufau G. Metabolic Effects of Bile Acids in the Gut in Health and Disease. *Curr Med Chem* 2014;21:2822-2829.

48. Cariou B, van Harmelen K, Duran-Sandoval D, et al. Transient impairment of the adaptive response to fasting in FXR-deficient mice. *FEBS Lett* 2005;579:4076-4080.
49. Ma K, Saha PK, Chan L, Moore DD. Farnesoid X receptor is essential for normal glucose homeostasis. *J Clin Invest* 2006;116:1102-1109.
50. Cariou B, van HK, Duran-Sandoval D, et al. The farnesoid X receptor modulates adiposity and peripheral insulin sensitivity in mice. *J Biol Chem* 2006;281:11039-11049.
51. Zhang Y, Lee FY, Barrera G, et al. Activation of the nuclear receptor FXR improves hyperglycemia and hyperlipidemia in diabetic mice. *Proc Natl Acad Sci U S A* 2006;103:1006-1011.
52. Sinal CJ, Tohkin M, Miyata M, Ward JM, Lambert G, Gonzalez FJ. Targeted disruption of the nuclear receptor FXR/BAR impairs bile acid and lipid homeostasis. *Cell* 2000;102:731-744.
53. Lambert G, Amar MJ, Guo G, Brewer HB, Jr., Gonzalez FJ, Sinal CJ. The farnesoid X-receptor is an essential regulator of cholesterol homeostasis. *J Biol Chem* 2003;278:2563-2570.
54. Watanabe M, Houten SM, Matakai C, et al. Bile acids induce energy expenditure by promoting intracellular thyroid hormone activation. *Nature* 2006;439:484-489.
55. Angelin B, Einarsson K, Hellstrom K, Leijd B. Effects of cholestyramine and chenodeoxycholic acid on the metabolism of endogenous triglyceride in hyperlipoproteinemia. *J Lipid Res* 1978;19:1017-1024.
56. Beil U, Crouse JR, Einarsson K, Grundy SM. Effects of interruption of the enterohepatic circulation of bile acids on the transport of very low density-lipoprotein triglycerides. *Metabolism* 1982;31:438-444.
57. Bateson MC, Maclean D, Evans JR, Bouchier IA. Chenodeoxycholic acid therapy for hypertriglyceridaemia in men. *Br J Clin Pharmacol* 1978;5:249-254.
58. Carulli N, Ponz de LM, Podda M, et al. Chenodeoxycholic acid and ursodeoxycholic acid effects in endogenous hypertriglyceridemias. A controlled double-blind trial. *J Clin Pharmacol* 1981;21:436-442.
59. Hambruch E, Miyazaki-Anzai S, Hahn U, et al. Synthetic farnesoid X receptor agonists induce high-density lipoprotein-mediated transhepatic cholesterol efflux in mice and monkeys and prevent atherosclerosis in cholesteryl ester transfer protein transgenic low-density lipoprotein receptor (-/-) mice. *J Pharmacol Exp Ther* 2012;343:556-567.
60. Kawamata Y, Fujii R, Hosoya M, et al. A G protein-coupled receptor responsive to bile acids. *J Biol Chem* 2003;278:9435-9440.
61. Maruyama T, Miyamoto Y, Nakamura T, et al. Identification of membrane-type receptor for bile acids (M-BAR). *Biochem Biophys Res Commun* 2002;298:714-719.
62. Keitel V, Haussinger D. Perspective: TGR5 (Gpbar-1) in liver physiology and disease. *Clin Res Hepatol Gastroenterol* 2012;36:412-419.
63. Pols TW, Nomura M, Harach T, et al. TGR5 activation inhibits atherosclerosis by reducing macrophage inflammation and lipid loading. *Cell Metab* 2011;14:747-757.